

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
**Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

**THIS PAGE BLANK (USPTO)**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : A23L 1/317, A23K 1/10, 1/18, A23J 3/22		A1	(11) International Publication Number: WO 97/02760
			(43) International Publication Date: 30 January 1997 (30.01.97)
(21) International Application Number: PCT/EP96/03060			(81) Designated States: AU, BR, CA, CN, CZ, FI, HU, JP, KR, MX, NO, NZ, PL, SG, TR, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(22) International Filing Date: 12 July 1996 (12.07.96)			
(30) Priority Data: 60/001,075 12 July 1995 (12.07.95) US			
(71) Applicant: SOCIETE DES PRODUITS NESTLE S.A. [CH/CH]; P.O. Box 353, CH-1800 Vevey (CH).			
(72) Inventors: POPPEL, Gerhard, J.; 512 Raintree Drive, St. Joseph, MO 64506 (US). RAYNER, Michael, G.; 9955 State Highway FF, Agency, MO 64401 (US). SAYLOCK, Michael, J.; 3621 N.W. 75th Court, Kansas City, MO 64151 (US).			
(74) Agent: McCONNELL, Bruce; Société des Produits Nestlé S.A., P.O. Box 353, CH-1800 Vevey (CH).			<b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: FORMULATED EMULSION PRODUCT AND PROCESS			
(57) Abstract			
<p>A process for the production of a formulated emulsion product which has a meat-like appearance. A protein source is emulsified and then alkaline is added to the emulsion to raise the pH to above about 8. The emulsion is rapidly treated in an emulsion mill to cause the protein in the heated emulsion to at least partially coagulate. The heated emulsion is then allowed to coagulate and form striations and is cut into chunks. The formulated emulsion product so produced contains about 45 % to 85 % by weight moisture and has a chewy texture and a high striated appearance.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Formulated Emulsion Product and Process

This invention relates to a process for the production of a formulated emulsion ~~product which has a meat-like appearance~~. The invention also relates to products so produced. The formulated emulsion products are particularly suitable for use as pet foods.

Coagulated meat emulsions, in the form of loaf-like products, are commonly used as pet foods because they are easily manufactured, readily digested by the animals, very palatable to the animals, and are readily formulated to contain necessary nutrients and trace elements. A disadvantage is that these coagulated emulsions do not have a striated and chunky meat-like appearance. However, for pet foods, a meat-like appearance can greatly enhance consumer acceptability.

A formulated meat emulsion which has a meat-like appearance is described in US patent 4,781,939. This formulated meat emulsion is produced by first forming a meat emulsion from a meat source. Dry ingredients such as dry proteinaceous materials (for example wheat gluten, soy flour), vitamins, minerals and the like are then mixed into the meat emulsion to provide a viscous emulsion. The viscous emulsion is then run through a high-speed emulsion mill in which the emulsion is rapidly heated to a temperature in the range of 102°C to 118°C. The emulsion leaving the emulsion mill is fed to a holding tube where the protein in the emulsion coagulates to form a solid emulsion product. This solid emulsion product is then formed into chunks. The chunks are highly striated and resemble natural meat chunks in appearance and texture.

The product produced by the process has been successfully used in pet foods for many years. However in recent times, mainly to reduce costs, there has been a move to increase the moisture content of the meat emulsions. However, if the amount of moisture becomes too high, the emulsion is not viscous enough to form acceptable chunks. Consequently proteinaceous materials must be added to the emulsion prior to forming into chunks.

A similar process is disclosed in US patent 5,132,137. However, in this process the viscous emulsion is heated to a temperature of 40 to 70°C in the emulsion mill; which is much lower than that in the process disclosed in US patent 4,781,939. The heated emulsion takes longer to coagulate and is therefore held in a holding tube for a longer time. The emulsion is then formed into strands and baked in an oven at a core temperature of 70 to 95°C. The patent is deficient

in specific teaching of the contents of the emulsion but it is very likely that proteinaceous components such as gluten would be needed and that the moisture content would need to be low. This is particularly so since only partial coagulation occurs in the emulsion mill.

5 Therefore there is a need for a process of producing a formulated emulsion product which has a meat-like appearance and texture and yet which may have a higher moisture content.

In one aspect, this invention provides a process for the production of a formulated emulsion product which has a meat-like appearance, the process  
10 comprising:

emulsifying a protein source for providing an emulsion;  
adding alkaline to the emulsion to raise the pH to above about 8;  
rapidly heating and comminuting the emulsion using mechanical energy  
for causing protein in the heated emulsion to at least partially coagulate;  
15 allowing the heated emulsion to coagulate and form striations; and  
forming the coagulated emulsion into chunks for providing a formulated emulsion product.

Surprisingly, it is found that the addition of the alkaline greatly increases the viscosity of the emulsion and permits the production of a formulated  
20 emulsion product of higher moisture content. Also, if desired, components such as gluten may be reduced in amount or omitted entirely. Further, ~~the emulsion need not be produced from a traditional meat source; instead an animal protein source such as plasma, blood proteins, whey, or casein may be used~~  
Consequently, a formulated emulsion product with a meat-like appearance may  
25 be produced at lower cost.

Further, it is also surprisingly found that the formulated emulsion product has much thinner and more pronounced layers giving it a highly striated and more meat-like appearance. Also the product has an improved, chewy texture.

The protein source may be of animal or vegetable origin, or a mixture of  
30 both. Conveniently the protein source may comprise at least about 80% by weight of animal protein; the remaining protein being of vegetable origin. For example, the protein in the emulsion may comprise about 95% to 100% by weight of protein of animal origin. Alternatively, the protein in the emulsion may comprise about 95% to 100% by weight of protein of vegetable origin.

35 The alkaline which is added to the emulsion may be any suitable alkaline which is acceptable in food. For example, the alkaline may be NaOH, KOH or

Ca(OH)<sub>2</sub>, or mixtures thereof. Preferably, however, the alkaline is NaOH. Preferably sufficient alkaline is added to raise the pH of the emulsion to a range of about 9 to about 12; for example about 9 to about 11. This amount preferably results in the formulated emulsion product containing less than about 5% by weight alkaline; more preferably less than about 2% by weight.

The formulated emulsion product ~~preferably has a moisture content of about 45% to about 85% by weight;~~ more preferably about 55 to about 65% by weight. If necessary water may be added up to an amount of about 60% by weight to obtain this level of moisture. However, if the protein source contains sufficient moisture, the addition of further moisture is not necessary. Additional ingredients such as fats, sugars, salts, spices, seasonings, flavorants, minerals, and the like may also be added to the emulsion. The amount of additional ingredients used is preferably such that they make up about 1% to about 35% by weight of the formulated emulsion product.

The protein source may include whey. This embodiment has the advantage that sugars in the whey may react with amino acids in the emulsion. The Maillard reaction products which result provide the chunks with "meaty" flavor notes which may increase palatability.

The alkaline may be added to the emulsion during emulsification of the protein source, or may be added to the emulsion immediately prior to the heating and comminuting of the emulsion, or may be added to the emulsion during the heating and comminuting of the emulsion. Preferably, the emulsion is deaerated after emulsification.

Preferably the emulsion is rapidly heated and comminuted by forcing the emulsion between a pair of spaced-apart plates; at least one of which is rotating at high speed. If necessary, the emulsion may be further heated by injecting superheated steam into it. Preferably the emulsion is heated to a temperature above about 100°C, for example within the range of about 102°C to about 120°C. Alternatively, the emulsion may be rapidly heated to a lower temperature providing that the temperature is sufficient to cause at least partial coagulation of the protein; for example at least to about 45°C.

The heated emulsion is preferably allowed to coagulate and form striations for a time in the range of about 30 seconds to about 10 minutes; more preferably about 1 minute to about 6 minutes.

The invention also provides a formulated emulsion product produced by the process defined above.

~~In another aspect, this invention provides a formulated, striated emulsion product having a meat-like appearance, the product comprising protein, fats and about 45% to 85% by weight moisture and having a pH of above about 8.~~

5 Preferably the formulated emulsion product comprises about 55% to about 65% moisture. Also, the formulated emulsion product preferably has a pH in the range of about 9 to about 12.

The protein content of the formulated emulsion product is preferably about 10 to about 25 % by weight; for example about 12 to about 22 % by weight. The protein may include whey.

10 The formulated emulsion product preferably comprises about 5% to about 25% by weight of fat.

15 Preferably at least about 50% of the striations of the formulated emulsion product have a thickness less than about 50 $\mu$ m. Further, the formulated emulsion product preferably has an elasticity recovery of greater than about 40%; more preferably greater than about 50%. For example, the formulated emulsion product may have an elasticity recovery of about 55% to about 65%. This provides the product with an excellent chewy texture.

20 The invention also provides a pet food comprising chunks of a formulated emulsion product as defined above, in combination with an acidic carrier to neutralize the basicity of the chunks. The carrier may be an acidic sauce or gravy, or may be a coagulated meat emulsion.

Embodiments of the invention are now described, by way of example only, with reference to the drawing which illustrates schematically the process of producing the formulated emulsion product.

25 Referring to the drawing, a protein source 2, water 4 and an additive mixture 6 are mixed together in a mixer 8 to provide a primary emulsion 10. The protein source 2 which is fed to the mixer 8 may be any suitable meat or vegetable protein source, or both.

30 A suitable meat protein source is a meat emulsion obtained by grinding and then emulsifying blocks of a meat material, such as meat or meat by-products. The meat material may be any suitable source of animal protein; for example the muscular or skeletal meat of mammals, poultry, and fish or meat by-products such as hearts, liver, kidneys, tongue and the like. The exact composition may be selected according to cost and the desired flavor. The meat material conveniently may be in frozen form prior to grinding. Suitable  
35 procedures for processing frozen blocks of meat material into a meat emulsion



mixture are described in US patent 4,781,939, the disclosure of which is incorporated by reference. Alternatively or in addition, ~~the animal protein source may be in the form of~~ meat meals such as poultry meal, fish meal, and red meat meal; ~~blood proteins~~; egg solids; casein; plasma; whey and mixtures thereof.

- 5 Again the exact composition may be selected according to cost and the desired flavor. Mixtures of any of the above may also be used.

Any suitable vegetable protein source may be used; for example grain flours and legume flours, vegetable protein isolates and concentrates, and vegetable proteins. Suitable examples are wheat flours, cottonseed meal, gluten, 10 soy meal, soy protein concentrates and isolates, and mixtures thereof. Preferably however the vegetable protein source contains reduced amounts of starch; for example, less than about 20% by weight of starch. Cottonseed meal, gluten, soy meal, soy protein concentrates and isolates are therefore particularly preferred.

- It is particularly preferred to use a protein source which is a mixture of 15 animal and vegetable origin; for example a mixture of meat material, soy protein isolates or concentrates, and gluten. The total amount of protein in the primary emulsion must be such that the protein is able to coagulate to provide a firm emulsion product in a relatively short time; for example in less than 10 minutes. For example, the primary emulsion 10 may contain from 10 to 22 % by weight of 20 protein.

- If desired, whey may be included in the protein source 2; for example in an amount of about 1% to about 25% by weight of the primary emulsion 10. The whey is conveniently in powder form. The inclusion of the whey has the advantage of providing a source of reducing sugars which may react with amino 25 acids to provide Maillard reaction products. These reaction products may provide flavor notes which can increase the palatability of the formulated emulsion product.

- The additive mixture 6 may be a mixture of dry additives or fats; and mixtures thereof. The dry additives may be, for example, sugar, salt, spices, 30 seasonings, vitamins, minerals, flavorants and the like. The fats may be suitable animal fats; for example tallow, or may be vegetable fats. The total amount of the additive mixture used may be selected as required but will usually be in the range of 1 to 35% by weight of the primary emulsion.

- The amount of water 4 added is sufficient such that the water comprises 35 from about 45% to 85% by weight in the formulated emulsion product. If sufficient moisture is present in the protein source 2 or the additive mixture 6, the

water 4 may be omitted. Usually an amount of water to make up to about 60% of the primary emulsion 10 is added.

In the processes described in US patents 4,781,939 and 5,132,137, the amount of fat in the primary emulsion must be controlled to facilitate processing and to obtain an acceptable product. However the amount of fat is not an important parameter in this process and the amount can be selected as desired. Consequently no fat need be added; particularly if the animal protein source 2 contains fats. Conveniently, the amount of fat in the primary emulsion 10 is in the range of 5% to 25% by weight. Reducing the importance of fat as a parameter in the process is a major advantage of the process.

The protein source 2, the water 4 and the additive mixture 6 are mixed in any suitable mixer 8. For example, a twin screw mixer, a twin ribbon blender, or an overlapping paddle mixer may be used. The mixing should be sufficient to ensure that any dry ingredients are dissolved.

Once mixed, the primary emulsion 10 is preferably fed through a vacuum stuffer (not shown), or similar deaeration apparatus, to deaerate the primary emulsion 10. This removes air which may otherwise cause disruption of the formulated emulsion product and reduce its meat-like appearance. However, it is not essential to deaerate the primary emulsion 10 because, in general, an acceptable product may still be obtained.

Alkaline 12 is injected into the primary emulsion 10 prior to the primary emulsion 10 being fed into an emulsion mill 14. Sufficient alkaline 12 is injected so as to raise the pH of the primary emulsion 10 to a range of about 8 to about 12. The alkaline may be any alkaline suitable for use in a food product.

Conveniently, NaOH may be used; preferably in solution. The concentration of the NaOH in solution is conveniently about 25% to 50% by weight. As an alternative to injecting the alkaline into the primary emulsion 10, the alkaline may be added into the mixer 8 along with the water 4 and additive mixture 6. As a further alternative, the alkaline may be injected into the primary emulsion 10 as it enters the emulsion mill 14.

The emulsion mill 14 is a high speed mill which subjects the emulsion to rapid mechanical heating and shearing. Any suitable emulsion mill 14 may be used, for example the emulsion mill disclosed in US patent 5,132,137, the disclosure of which is incorporated by reference. Other suitable emulsion mills 14 are commercially available under the tradename of Trigonal and may be obtained from Siefer Maschinenfabrik GmbH & Co KG, Bahnhofstrasse 114,

Postfach 101008, Velbert 1, Germany. These emulsion mills usually comprise a pair of plates which are closely spaced apart. One of the plates rotates at high speeds. The emulsion is fed between the plates and the mechanical energy imparted to the emulsion rapidly raises its temperature. The heated emulsion 16 leaves the emulsion mill in a thin stream which packs onto earlier streams to form layers or striations

The temperature of the primary emulsion 10 is raised to the desired coagulation temperature in the emulsion mill 14 in a few seconds. For example, the temperature may be raised to from about 100°C to about 120°C. Alternatively, the temperature may be raised to in the range of about 45°C to about 75°C as described in US patent 5,132,137. Usually the mechanical energy generated in the emulsion mill 14 will be sufficient to heat the emulsion 10 but this may be supplemented by the injection of superheated steam.

The heated emulsion 16 leaving the emulsion mill 14 is then transferred to a holding tube 18. Holding tubes 18 such as described in US patents 4,781,939 and 5,132,137 may be used. In the holding tube 18, the heated emulsion 16 coagulates while moving slowly along the holding tube 18. The residence time of the heated emulsion 16 in the holding tube 18 is sufficient for the emulsion to have coagulated into a firm emulsion product upon reaching the exit of the holding tube 18. Residence times of about 30 seconds to about 10 minutes are sufficient; the actual time depending upon the temperature to which the primary emulsion 10 was heated. Longer or shorter residence times may also be used but disadvantages arise. If desired, the holding tube may be equipped with a suitable valve at its end to maintain a desired pressure in it.

The firm emulsion product 20 leaving the holding tube 18 is then transferred to a cutter 22 where it is cut into chunks 24 of size suitable for use in a pet food. Alternatively, if it is desired to produce chunks of a more natural appearance, steam may be intermittently injected into the holding tube 18 to cause disruption of the firm emulsion product 20 into chunks as it leaves the holding tube 18. This procedure is described in US patent 4,781,939.

If desired, the chunks 24 may be baked by passing them through an oven as described in US patent 5,132,137. However, if the meat emulsion 10 was heated to above 100°C in the emulsion mill 14, this is unnecessary.

The chunks 24 are then allowed to cool and are collected. If desired, the collected chunks are stored for later use or are simply transferred to a canning operation. Prior to canning, the chunks 24 may be flaked if desired. In the

canning operation, the chunks are filled into cans along with a gravy or sauce. The gravy or sauce may contain an acid, such as citric acid or phosphoric acid, which is acceptable in foods to neutralize the alkaline in the chunks. The cans are then sealed and sterilized. In this case, the product produced is a chunk-in-gravy type product. Alternatively the chunks may be combined with a meat emulsion which forms a loaf-type product. In this case, the meat emulsion may contain a similar acid to neutralize the chunks. The cans are then sealed and sterilized. In this case, the product produced is a chunk-in-loaf type product.

5 The chunks 24 have a highly striated appearance which provides a very good simulation of meat. This greatly increases the consumer acceptability of the product. Also, the chunks 24 have a chewy texture which requires the animals to chew them; much as animals would chew fresh meat. This offers the advantage of being beneficial to the animals' teeth.

10 Although the invention has been described with reference to pet foods, it will be appreciated that the process may also be used to produce foods intended for human consumption.

#### Example 1

20 An emulsion is formed by mixing 20% plasma, 20% whey powder, 6% tallow, and 52% water (the percentages being based on the total weight of the chunk ingredients). The emulsion is transferred to a batch blender and 2% of a solution containing 50% by weight of NaOH is added. The pH of the emulsion increases to about 11.

25 The emulsion is run through an emulsion mill (a Trigonal Mill obtained from Siefer Maschinenfabrik GmbH & Co KG). The emulsion leaving the emulsion mill is at a temperature of 103 to 105°C and is discharged into a holding tube. The residence time in the holding tube is less than 6 minutes. The coagulated emulsion leaving the holding tube is cut into chunks of about 10 mm size. The chunks have a highly striated, meat-like appearance. Some meat-like browning, due to Maillard reactions between amino acids in the plasma and sugars in the whey, is noticeable.

30 A loaf product mix is prepared from about 53% by weight of meat and meat by-products, about 44.5% water and about 2.5% flavoring agents. The mixture is cooked in a batch mixer/cooker and filled, with the chunks, into cans.

The mass ratio of loaf ingredients to chunks is about 78:22. The cans are sealed and sterilized.

A can is opened, a representative sample of the product is taken and the pH is determined to be about 6.5 to 7.0. Further product is fed to dogs and is  
5 freely eaten.

### Example 2

Chunks are produced as described in example 1. A gravy is then prepared  
10 by mixing together 5.5% starch, 1% phosphoric acid, less than 0.5% condiments, and water making up the remainder (all percentages being by weight). The chunks and gravy are combined in a mass ratio of about 45:55 and filled into cans. The cans are sealed and sterilized.

A can is opened, a representative sample of the product is taken and the  
15 pH is determined to be about 6.3 to 6.6. Further product is fed to dogs and is freely eaten.

### Example 3

20 Frozen meat and meat by-products obtained from a local source are ground and then emulsified. An emulsion mixture is then formed by mixing ~~2%~~  
~~plasma~~, 2% whey powder, 1% condiments and 93% emulsified meat and meat by-products (the percentages being based on the total weight of the chunk ingredients). The emulsion mixture is transferred to a batch blender and 2% of a  
25 solution containing 50% by weight of NaOH is added. The pH of the emulsion mixture increases to about 11.

The emulsion mixture is run through an emulsion mill (a Trigonal Mill obtained from Siefer Maschinenfabrik GmbH & Co KG). The emulsion leaving the emulsion mill is at a temperature of 103 to 105°C and is discharged into a  
30 holding tube. The residence time in the holding tube is less than 6 minutes. The coagulated emulsion leaving the holding tube is cut into chunks of about 10 mm size. The chunks have a highly striated, meat-like appearance. Some meat-like browning, due to Maillard reactions between amino acids in the plasma and sugars in the whey, is noticeable.

35 A gravy is then prepared by mixing together 5.5% starch, 1% phosphoric acid, less than 0.5% condiments, and water; the water making up the remainder

-10-

(all percentages being by weight). The chunks and gravy are combined in a mass ratio of about 45:55 and filled into cans. The cans are sealed and sterilized.

A can is opened, a representative sample of the product is taken and the pH is determined to be about 6.3 to 6.6. Further product is fed to cats and dogs  
5 and is freely eaten.

#### Example 4

Frozen meat and meat by-products obtained from a local source are  
10 ground and then emulsified. The emulsion is placed in a blender and 10% by weight wheat gluten, 4% by weight soy protein concentrate, 3% condiments and 18 % by weight water are added to provide an emulsion mixture (the percentages being based on the total weight of all chunk ingredients). Two % by weight of a solution containing 50% by weight of NaOH is added then added and the  
15 emulsion mixture blended until uniform. The pH of the emulsion mixture increases to about 12.

The emulsion mixture is run through a Trigonal emulsion mill and discharged into a holding tube at a temperature of about 110°C to about 115°C. The holding tube is equipped with a pinch valve to maintain back pressure. The  
20 coagulated emulsion leaving the holding tube is cut into chunks of about 10 mm size. The chunks have a shredded, stringy, meat-like appearance.

A gravy is then prepared by mixing together 2.5% starch, 0.5% condiments, 0.3% hydrocolloid, and water; the water making up the remainder (all percentages being by weight on the basis of gravy ingredients). The chunks  
25 and gravy are combined in a mass ratio of about 60:40 and then about 1.75% by weight of phosphoric acid is added and blended in. The mixture is then filled into cans. The cans are sealed and sterilized.

#### Example 5

30 An emulsion is prepared by blending 8% by weight wheat gluten, 22% by weight soy protein concentrate, 0.3% condiments, 5% fat and 62.7 % by weight water (the percentages being based on the total weight of all chunk ingredients). Two % by weight of a solution containing 50% by weight of NaOH is added then  
35 added and the emulsion mixture blended until uniform. The pH of the emulsion mixture increases to about 11.

The emulsion is then formulated into chunks and added to a gravy as described in example 4.

#### Example 6

5

A chunk produced according to the process described in US patent 4,781,939 (Sample A) and a chunk produced according to example 4 (Sample 1) are frozen at -80°C. The samples are freeze fractured into sections of 10 to 15  $\mu\text{m}$  and stained with Sudan III and Harris' Hematoxylin (*J. Dairy Science*; 1922, 10 5). The samples are then subjected to light microscopy in the frozen state.

Sample 1 is formed of thin layers or striations of protein and fat generally of thickness less than about 50  $\mu\text{m}$ . Sample A is formed of less well defined layers or striations of protein and fat of thickness substantially generally greater than 50  $\mu\text{m}$ . The results support the visual observation that the chunks of Sample 1 have an appearance more highly striated then the conventional chunks of 15 Sample A.

#### Example 7

20

A dough which contains about 62.5% by weight of a mixture of turkey meat and fish is prepared according to example 4 (without the formation of chunks). The dough are labeled Sample 2. A similar dough is produced according to the process described in US patent 4,781,9394 (again without the formation of chunks). The dough is labelled Sample B. The doughs are then cut 25 into disks of about 50mm diameter and 10mm thickness. Twenty disks of sample 2 are prepared in this way and 20 disks of sample B.

Each sample disk is placed in a texture analyzer (TX.XT2 by Texture Technologist Corp, Scarsdale, New York, USA) on a flat disk of 80 mm. The temperature is about 10 to 15°C. A flat bottom probe of 56 mm diameter is 30 pressed down at a constant speed of 2 mm/s on the sample disk using 10 gram force and for a distance of 6 mm. The probe is held in this position for 60 seconds. The strain at the start of the pressing (initial strain) and the strain after pressing (final strain) are determined and recorded. The percentage elasticity recovery of each disk is then determined according to the formula:

35

$$\text{recovery \%} = 100 - ((\text{final strain} - \text{initial strain})/\text{final strain} \times 100)$$

The results are as follows:

Sample 2				Sample B			
Disk	Final strain	Initial strain	% Recovery	Disk	Final strain	Initial strain	% Recovery
1	1402	887	63.3	1	812	223	27.5
2	1475	943	64.0	2	763	208	27.2
3	838	519	62.0	3	543	136	25.0
4	443	272	61.4	4	829	233	28.1
5	308	196	63.8	5	810	213	26.3
6	610	389	63.7	6	1011	288	28.5
7	930	583	62.7	7	781	216	27.6
8	953	579	60.7	8	616	168	27.3
9	493	309	62.7	9	799	200	25.1
10	501	314	62.7	10	771	217	28.2
11	462	285	61.7	11	755	207	27.4
12	356	222	62.0	12	509	140	27.5
13	683	429	62.7	13	821	212	25.8
14	425	260	61.2	14	532	141	26.6
15	492	307	62.4	15	672	181	27.0
16	87	53	60.7	16	538	151	28.0
17	801	489	61.1	17	843	229	27.2
18	1253	771	61.5	18	695	180	25.8
19	911	567	62.2	19	682	184	27.0
20	325	194	59.6	20	524	142	27.0

5

The average % recovery for the disks of sample 2 is 63.3 while that for the disks of sample B is 27.0. This indicates that chunks produced from the dough of Sample 2 have a much more elastic (chewier) texture than the conventional chunks.



Claims

1. A process for the production of a formulated emulsion product which has a meat-like appearance, the process comprising:
  - 5 emulsifying a protein source for providing an emulsion;
  - adding alkaline to the emulsion to raise the pH to above about 8;
  - rapidly heating and comminuting the emulsion using mechanical energy for causing protein in the heated emulsion to at least partially coagulate;
  - allowing the heated emulsion to coagulate and form striations; and
  - 10 forming the coagulated emulsion into chunks for providing a formulated emulsion product.
2. A process according to claim 1 in which NaOH, KOH or Ca(OH)<sub>2</sub>, or mixtures thereof, is added as the alkaline.
- 15 3. A process according to claim 2 in which sufficient alkaline is added to raise the pH of the emulsion to a range of about 9 to about 12.
4. A process according to claim 1 in which up to about 60% by weight, on  
20 the basis of the weight of the emulsion, of moisture is emulsified with the protein source for providing a formulated emulsion product having a moisture content of about 45% to 85% by weight.
5. A process according to claim 1 in which the emulsion is heated to a  
25 temperature in the range of about 102°C to about 120°C.
6. A formulated, striated emulsion product having a meat-like appearance, the product comprising protein, fats and about 45% to 85% by weight moisture and having a pH of above about 8.
- 30 7. A formulated emulsion product according to claim 6 which comprises about 55% to about 65% by weight moisture.
8. A formulated emulsion product according to claim 6 in which the protein  
35 comprises about 10% to about 25% by weight of the product.

9. A formulated emulsion product according to claim 6 in which at least about 50% of the striations have a thickness of less than about 50 $\mu$ m.
10. A pet food comprising chunks of a formulated emulsion product according to any of claims 6 to 9, in combination with an acidic carrier to neutralize the basicity of the chunks.




Fig. 1

# INTERNATIONAL SEARCH REPORT

Inter. Application No  
PCT/EP 96/03060

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A23L1/317 A23K1/10 A23K1/18 A23J3/22

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A23L A23K A23J

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US,A,4 781 939 (R. CRAIG MARTIN ET AL.) 1 November 1988 cited in the application see the whole document	1,4-8
Y	WO,A,92 18018 (NADREPH LIMITED) 29 October 1992 see page 2, paragraph 2 - paragraph 4 see page 3, paragraph 2 - paragraph 3 see page 5, paragraph 2 - paragraph 3 see example 7 see claims 1,10,14,17-20	1,4-8
A	---	2-4
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

7 November 1996

Date of mailing of the international search report

20. 11. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Dekeirel, M

## INTERNATIONAL SEARCH REPORT

Inter national Application No  
PCT/EP 96/03060

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE WPI Week 7804 Derwent Publications Ltd., London, GB; AN 78-07553A XP002017903 & JP,A,52 148 652 (IKAWA M) , 10 December 1977 see abstract ---	1
A	US,A,3 843 815 (STANLEY H. REESMAN) 22 October 1974 see claim 1 ---	10
A	US,A,5 132 137 (JOSEF REIMANN ET AL.) 21 July 1992 cited in the application see the whole document ---	1,6
A	EP,A,0 651 948 (SOCIETE DES PRODUITS NESTLE S.A.) 10 May 1995 see the whole document -----	1

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 96/03060

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4781939	01-11-88	CA-A- 1328051	29-03-94
		AU-B- 596048	12-04-90
		AU-A- 7929487	05-05-88
		DE-A- 3779397	02-07-92
		EP-A- 0265740	04-05-88
		FI-C- 91034	10-05-94
		IE-B- 60400	13-07-94
		JP-A- 63123356	27-05-88
		ZA-A- 8707544	13-04-88
WO-A-9218018	29-10-92	AT-T- 143769	15-10-96
		AU-B- 657754	23-03-95
		AU-A- 1648292	17-11-92
		CA-A- 2104863	23-10-92
		EP-A- 0581814	09-02-94
		JP-T- 6506589	28-07-94
		US-A- 5480973	02-01-96
US-A-3843815	22-10-74	NONE	
US-A-5132137	21-07-92	DE-C- 3933528	06-12-90
		AU-B- 635343	18-03-93
		AU-A- 6416290	28-04-91
		CA-A- 2035476	05-04-91
		WO-A- 9104677	18-04-91
		EP-A- 0422750	17-04-91
		JP-B- 6067306	31-08-94
		JP-T- 4504062	23-07-92
EP-A-651948	10-05-95	AU-A- 7428894	27-04-95
		CA-A- 2133502	12-04-95
		JP-A- 7184600	25-07-95
		NO-A- 943699	12-04-95
		ZA-A- 9407577	15-05-95